

# **KINETIC AND THERMODYNAMIC STUDIES OF THE LIGAND SUBSTITUTION REACTIONS OF THE COBALAMINS**

**Leanne Knapton**

*A thesis submitted to the Faculty of Science,  
University of the Witwatersrand,  
Johannesburg,  
In fulfilment for the requirement for the  
Degree of Doctor of Philosophy*

*March 2005*

**DECLARATION**

I declare that this is my own unaided work. It is being submitted for the Degree of Doctor of Philosophy at the University of the Witwatersrand, Johannesburg. It has not been submitted before for any degree or examination at any other university.

---

L. Knapton

31 March 2005

**ABSTRACT**

The ligand substitution reactions of aquacobalamin are fast and hence the usual inertness of the  $d^6$  Co(III) ion has been modified. It is well established that the reactions proceed through a dissociative interchange mechanism; however, previous ligand studies were performed in a KCl medium, which led to the formation of the more substitution-inert chloro complex. The kinetics of aquacobalamin were reinvestigated with the ligands  $N_3^-$ ,  $NO_2^-$ ,  $SCN^-$ ,  $S_2O_3^{2-}$ ,  $OCN^-$  and  $SeCN^-$  in a  $NaNO_3$  medium. The reactions proceeded too rapidly for saturation kinetics to be observed and hence only the second-order rate constants could be obtained. These were corrected for pH and determined as a function of temperature, from which the activation parameters were determined. The donor atom of the ambidentate ligands were investigated and correlations were found between the Mulliken population on the donor atom, the energy of the highest occupied molecular orbital (HOMO) with  $\sigma$  symmetry, and  $\Delta H_{k_{II}}^\ddagger$ , the enthalpy of activation, and  $\Delta S_{k_{II}}^\ddagger$ , the entropy of activation, respectively. Good correlations occurred when the donor atoms were taken to be N for  $SCN^-$  and  $NO_2^-$ ; S for  $S_2O_3^{2-}$ ; O for  $OCN^-$  and Se for  $SeCN^-$ .

The effect that changing the environment of aquacobalamin has on its kinetics was observed by determining the rate constants for the reaction of pyridine with aquacobalamin in water and 70% ethanol. The rates were faster in water and the activation parameters obtained for the reaction of aquacobalamin with pyridine in 70% ethanol are larger than they are for the reaction in water. The larger  $\Delta H^\ddagger$  arises due to less bond formation between pyridine and Co in the transition state and  $\Delta S^\ddagger$  is larger because it is dominated by the freeing of the coordinated water i.e. bond breaking is the dominant process in the transition state.

The effects of a bulkier ligand than water on the kinetics of aquacobalamin were investigated. The temperature dependence of the kinetics of the substitution of  $I^-$  in iodocobalamin by imidazole,  $N_3^-$  and  $S_2O_3^{2-}$  was studied. Despite the increase in size of the departing ligand there is still nucleophilic participation of the

incoming ligand in the transition state and hence the reaction still proceeds via an  $I_d$  mechanism.

In order to probe the *cis*-effect of the corrin in vitamin B<sub>12</sub> derivatives, comparative studies were undertaken of the reactions of aquacobalamin and aqua-10-Xcobalamin, X = Cl, NO, NH<sub>2</sub>, where the H at C10 was replaced with an electron-donating (Cl, NH<sub>2</sub>) or electron-withdrawing (NO) group. Formation constants were obtained for aquacobalamin and aqua-10-chlorocobalamin for the substitution of coordinated H<sub>2</sub>O with various anions (N<sub>3</sub><sup>-</sup>, NO<sub>2</sub><sup>-</sup>, SCN<sup>-</sup>, S<sub>2</sub>O<sub>3</sub><sup>2-</sup>, OCN<sup>-</sup>, SeCN<sup>-</sup>) and neutral *N*-donor ligands (CH<sub>3</sub>NH<sub>3</sub>, pyridine, imidazole). The anionic ligands bind more strongly to aqua-10-chlorocobalamin than to aquacobalamin with log *K* values larger by between 0.10 and 0.63 (average 0.26) larger. The converse is true for the neutral *N*-donor ligands, where log *K* is smaller by between 0.17 and 0.3 (average 0.25). Semi-empirical molecular orbital (SEMO) calculations using the ZINDO/1 model on the hydroxo complexes show that charge density is delocalised from the axial donor atom to the metal and Cl. Thus the anionic ligands bind more strongly to aqua-10-chlorocobalamin because of the ability of the metal and the Cl at C10 to accept charge density from the ligand. The cobalt ion in aqua-10-chlorocobalamin is more electron rich than it is in aquacobalamin and so it is less likely to accept further electron density from a neutral axial donor ligand. This results in the stability being lower than that of aquacobalamin.

The reaction kinetics of the substitution of H<sub>2</sub>O in aqua-10-chlorocobalamin were determined for the ligands N<sub>3</sub><sup>-</sup> and pyridine. The reaction proceeds via a dissociative interchange mechanism since saturation was seen for pyridine and not for N<sub>3</sub><sup>-</sup>. The activation parameters,  $\Delta H^\ddagger$  and  $\Delta S^\ddagger$ , are lower for aqua-10-chlorocobalamin than aquacobalamin and hence it can be deduced that bond breaking between the coordinated water and the cobalt atom is more dominant in aquacobalamin. The rates of reaction are faster for aquacobalamin than they are for aqua-10-chlorocobalamin. SEMO calculations show that as the Co–O bond is stretched, the charge density on Co in aquacobalamin is always lower than that on

aqua-10-chlorocobalamin, suggesting that aquacobalamin is a better electrophile towards the incoming ligand, thereby explaining the faster kinetics.

Aqua-10-nitrosocobalamin was synthesised and characterised by FAB(MS), NMR and UV-vis spectroscopy. The strongly electron-withdrawing NO group has deactivated the metal ion towards ligand substitution, with neither 1.2 M pyridine nor 0.7 M  $\text{N}_3^-$  showing any spectroscopic evidence for the displacement of the axial  $\text{H}_2\text{O}$  ligand. This provides further evidence that the electronic structure of the corrin ring can directly influence the ligand-binding properties of the metal.

Aqua-10-aminocobalamin was synthesised from aqua-10-nitrosocobalamin but is unstable in solution. Hence, only a preliminary UV-vis study could be undertaken with the compound. This study shows that the shifts in the bands occur towards longer wavelengths than that of aqua-10-chlorocobalamin, suggesting that the amino group at the C10 position donates more electron density to the cobalt centre than the chloro group.

*This work is dedicated to my Family:*

*My Parents: John and Deneen Knapton*

*Siblings: Cindy and Peter*

*Husband and soulmate: Mark*

*Dada and Mama – I wish you could have been here to see the end product.*

*Thanks to all of you for taking the journey with me and helping me to realise my  
dream.*

**ACKNOWLEDGEMENTS**

I would like to offer my sincerest thanks to the following people:

To my supervisor, Professor Helder Marques, for giving me the opportunity to be a part of the 'B<sub>12</sub> world';

Professor Kenneth Brown and Dr Xiang Zou at Ohio University in the USA for providing the aqua-10-chlorocobalamin, and collecting and interpreting the NMR spectroscopic data for aqua-10-nitrosocobalamin;

Basil Chassoulas, for plenty of help with the instruments;

Manuel Fernandes, for being as passionate about my crystals as I was and for all the late nights he spent trying to get something out of them.

A big thanks to the old gang from Lab 327 and 328; Alvaro de Sousa, Anton Lopic, Robby Luckay, Gary Patrick and Rachel Smith-Jasson for all their help and friendship over the years. Also, to all those from the Chemistry Department who have passed in and out of my life throughout this work, including Gaby Meirim and Sandy Lotz – thank you for making these the best years of my life.

A special thanks goes to my family – Dad and Mom – thanks for being the best parents in the world and giving me all the opportunities that you have given me; Cindy, Michael, Peter and Elmarie – thanks for all your love and support; Mark – you are the greatest, thanks for being a part of this, for listening to the moans when experiments were not working, and for the evenings and days that you kept me company in the lab; and lastly my best friends Manuel Fernandes, Chris Perry, Jose Oliveira and Preeti Vashi, for loads and loads of help, plenty of useful discussions and lots of coffee!

Thanks also go to the University of the Witwatersrand and the NRF for funding.

**LIST OF PUBLICATIONS**

1. H. M. Marques and L. Knapton, Factors affecting the rate of ligand substitution reactions of aquacobalamin (vitamin B<sub>12a</sub>), *Journal of the Chemical Society, Dalton Transactions*, 1997, 3827.
2. K. L. Brown, S. Cheng, X. Zou, J. S. Zubkowski, E. J. Valente, L. Knapton and H. M. Marques, *Cis* Effects in the Cobalt Corrins. 1. Crystal structures of 10-Chloroaquacobalamin perchlorate, 10-Chlorocyanocobalamin, and 10-Chloromethylcobalamin, *Inorganic Chemistry*, 1997, **36**, 3666.
3. H. M. Marques, L. Knapton, X. Zou and K. L. Brown, Probing the nature of the Co(III) ion in cobalamins: deactivation of the metal towards ligand substitution in 10-nitrosoaquacobalamin, and the kinetics of the ligand substitution reactions of iodocobalamin, *Journal of the Chemical Society, Dalton Transactions*, 2002, 3195.
4. L. Knapton and H. M. Marques, Probing the nature of the Co(III) ion in cobalamins: a comparison of the reaction of aquacobalamin (vitamin B<sub>12a</sub>) and aqua-10-chlorocobalamin with some anionic and *N*-donor ligands, *Dalton Transactions*, 2005, 889.



**CONTENTS**

	<b>Page</b>
<b>DECLARATION</b>	ii
<b>ABSTRACT</b>	iii
<b>DEDICATION</b>	vi
<b>ACKNOWLEDGEMENTS</b>	vii
<b>LIST OF PUBLICATIONS</b>	viii
<b>LIST OF FIGURES</b>	xvi
<b>LIST OF TABLES</b>	xxv
<b>LIST OF ABBREVIATIONS</b>	xxx
 <b>CHAPTER 1</b>	
<b>INTRODUCTION</b>	
<b>1.1 Vitamin B<sub>12</sub> and its Structure</b>	1
<b>1.2 Adenosylcobalamin (Coenzyme B<sub>12</sub>)</b>	6
<b>1.3 Cobaloximes</b>	9
<b>1.4 Porphyrin vs. Corrin</b>	10
<b>1.5 Coordination Chemistry of the Cobalamins</b>	11
<b>1.5.1 <i>cis</i>- and <i>trans</i>-Effects of the Cobalamins</b>	12
<b>1.5.2 Density Functional Theory (DFT) Calculations</b>	19
<b>1.6 Absorption Spectra of the Cobalamins</b>	21
<b>1.7 Ligand Substitution Reactions</b>	27
<b>1.7.1 Mechanism for the Ligand Substitution Reactions of the Cobalamins</b>	28
<b>1.8 Aims of this Study</b>	38
<b>References for Chapter 1</b>	40

**Contents (continued)****CHAPTER 2****EXPERIMENTAL**

<b>2.1</b>	Buffers	47
<b>2.2</b>	pH Measurements	47
<b>2.3</b>	Data Analysis	48
<b>2.4</b>	$pK_a$ Determinations	48
<b>2.5</b>	Kinetics of the Ligand Substitution Reactions of Aquacobalamin	49
<b>2.5.1</b>	Stopped-flow Spectrometry	49
<b>2.5.2</b>	Molecular Orbital Calculations	52
<b>2.6</b>	Solvent Effects	52
<b>2.7</b>	Kinetics of the Ligand Substitution Reactions of Iodobalamin	52
<b>2.8</b>	Formation Constants of Aquacobalamin and Aqua-10-chloro-cobalamin	53
<b>2.8.1</b>	Molecular Orbital Calculations	54
<b>2.9</b>	Ligand Substitution Reactions of Aqua-10-chlorocobalamin	54
<b>2.10</b>	Synthesis of Aqua-10-nitrosocobalamin	54
<b>2.10.1</b>	NMR Spectroscopy	56
<b>2.10.2</b>	FAB-Mass Spectrometry	56
<b>2.10.3</b>	Crystals of Aqua-10-nitrosocobalamin	56
<b>2.11</b>	Synthesis of Aqua-10-aminocobalamin	57
	References for Chapter 2	59

**CHAPTER 3** **$pK_a$  DETERMINATION OF AQUACOBALAMIN, AQUA-10-CHLOROCOBALAMIN, AQUA-10-NITROSOCOBALAMIN, AND AQUACOBALAMIN IN 70% ETHANOL**

<b>3.1</b>	Introduction	61
<b>3.2</b>	Results and Discussion	62
<b>3.2.1</b>	$pK_a$ Determination of Aquacobalamin	62

**Contents (continued)**

3.2.2	$pK_a$ Determination of Aqua-10-chlorocobalamin	64
3.2.3	$pK_a$ Determination of Aqua-10-nitrosocobalamin	66
3.2.4	$pK_a$ Determination of Aquacobalamin in 70% Ethanol	67
3.2.5	Molecular Orbital Calculations	69
3.2.6	Compensation Effects	70
3.3	Conclusion	71
	References for Chapter 3	72

**CHAPTER 4****LIGAND SUBSTITUTION REACTIONS OF AQUA-COBALAMIN**

4.1	Introduction	73
4.2	Results and Discussion	76
4.2.1	Second-Order Rate Constants for the Substitution of Water by Some Anionic Ligands	76
4.2.2	Compensation Effects Observed for Aquacobalamin	82
4.2.3	Coordination of the Ambidentate Ligands	90
4.2.3.1	A Comparison of the Predicted Donor Atoms with Those Reported in the Literature.	96
4.3	Conclusion	101
	References for Chapter 4	103

**CHAPTER 5****SOLVENT EFFECTS**

5.1	Introduction	105
5.2	Results and Discussion	106
5.3	Conclusion	116
	References for Chapter 5	117

**Contents (continued)****CHAPTER 6****THE KINETICS OF THE LIGAND SUBSTITUTION REACTIONS OF IODOCOBALAMIN**

<b>6.1</b>	Introduction	118
<b>6.2</b>	Results and Discussion	119
<b>6.2.1</b>	Substitution Kinetics of Iodocobalamin with the Imidazole and Azide Ligands	121
<b>6.2.2</b>	Substitution Kinetics of Iodocobalamin with the Thiosulfate Ligand	125
<b>6.2.2.1</b>	Determining the Outer-sphere Constant, $K_{OS}$ , for the Thiosulfate Ligand	126
<b>6.3</b>	Conclusion	132
	References for Chapter 6	133

**CHAPTER 7****MODIFICATION OF THE C10 POSITION OF AQUA-COBALAMIN**

	References for Chapter 7	141
--	--------------------------	-----

**CHAPTER 8****A COMPARISON OF THE FORMATION CONSTANTS OF COMPLEXES OF AQUACOBALAMIN AND AQUA-10-CHLOROCOBALAMIN**

<b>8.1</b>	Introduction	142
<b>8.2</b>	Results and Discussion	145
<b>8.2.1</b>	UV-vis Spectroscopic Study	145
<b>8.2.2</b>	Determination of Formation Constants	148
<b>8.2.2.1</b>	Semi-empirical Molecular Orbital Calculations	155
<b>8.3</b>	Conclusion	157

**Contents (continued)**

References for Chapter 8	159
--------------------------	-----

**CHAPTER 9****REACTION KINETICS FOR THE SUBSTITUTION OF  
COORDINATED H<sub>2</sub>O IN AQUA-10-CHLOROCOBALAMIN**

<b>9.1</b>	Introduction	161
<b>9.2</b>	Results and Discussion	163
<b>9.2.1</b>	Substitution Kinetics of Aquacobalamin and Aqua-10- chlorocobalamin with Pyridine	164
<b>9.2.2</b>	Substitution Kinetics of Aquacobalamin and Aqua-10- chlorocobalamin with Azide	169
<b>9.2.3</b>	Semi-empirical Molecular Orbital Calculations	177
<b>9.3</b>	Conclusion	183
	References for Chapter 9	184

**CHAPTER 10****AQUA-10-NITROSOCOBALAMIN**

<b>10.1</b>	Introduction	186
<b>10.2</b>	Results and Discussion	187
<b>10.2.1</b>	Ligand Binding Properties of Aqua-10-nitrosocobalamin	192
<b>10.2.2</b>	Crystals of Aqua-10-nitrosocobalamin	194
<b>10.3</b>	Conclusion	197
	References for Chapter 10	198

**CHAPTER 11****AQUA-10-AMINOCOBALAMIN**

<b>11.1</b>	Introduction	199
<b>11.2</b>	Results and Discussion	199

**Contents (continued)**

<b>11.3 Conclusion</b>	203
References for Chapter 11	204
 <b>CHAPTER 12</b>	
<b>CONCLUSIONS</b>	
References for Chapter 12	210
 <b>APPENDICES</b>	
<b>Appendix 1</b>	
NMR Spectroscopic Data for Aqua-10-nitrosocobalamin	I
<b>Appendix 2</b>	
FAB-Mass Spectrometry Spectra for Aqua-10-nitrosocobalamin	IX
<b>Appendix 3</b>	
Derivation of the Equations for a Dissociative and Dissociative Interchange Mechanism when $k_{-2}$ and $k_{-4}$ , Respectively, are Significant	X
Derivation of the Equations for a Dissociative and Dissociative Interchange Mechanism when $k_{-2}$ and $k_{-4}$ , Respectively, are Insignificant	XIV
<b>Appendix 4</b>	
Equation for the Binding of an Axial Ligand by Aquacobalamin and Aqua-10-chlorocobalamin	XVII
<b>Appendix 5</b>	
Determination of the Equations for the Correction of the Observed Rate Constants for pH and the Protonation of the Exogenous Ligand	XIX
<b>Appendix 6</b>	
Determination of the Equations for the Correction of the Observed Binding Constant for pH and the Protonation of the Exogenous Ligand	XXI
<b>Appendix 7</b>	
Calculation of the Outer-sphere Constant using the Fuoss-Eigen Equation	XXIII

**Contents** (*continued*)**Appendix 8**

Data for Crystal Structure of OCNcobalamin XXV

**Appendix 9**

Data Relevant to Chapter 3 CD

**Appendix 10**

Data Relevant to Chapter 4 CD

**Appendix 11**

Data Relevant to Chapter 5 CD

**Appendix 12**

Data Relevant to Chapter 6 CD

**Appendix 13**

Data Relevant to Chapter 8 CD

**Appendix 14**

Data Relevant to Chapter 9 CD

**Appendix 15**

Data Relevant to Chapter 10 CD

**Appendix 16**

Data Relevant to Chapter 11 CD

**LIST OF FIGURES**

<b>Figure</b>		<b>Page</b>
1.1	Family of uroporphinoid cofactors.	2
1.2	The structure and numbering scheme of the cobalamins where X = CN for cyanocobalamin and H <sub>2</sub> O for aquacobalamin.	3
1.3	Structure and numbering scheme of the adenosyl group, which forms the compound adenosylcobalamin when it is the X ligand in Figure 1.2. Attachment to the Co atom occurs through A15.	6
1.4	Coenzyme B <sub>12</sub> -dependent rearrangements.	7
1.5	Cobaloxime, a model compound for coenzyme B <sub>12</sub> where L and X can be a variety of ligands.	9
1.6	Correlation of the $\beta$ band with the C10 hydrogen chemical shift ( $R^2 = 0.97$ ).	16
1.7	Energy levels of the $\pi$ electron system and their corresponding bands in the absorption spectrum of aquacobalamin.	22
1.8	Molecular orbital diagram for [CH <sub>3</sub> Co <sup>III</sup> -(CR)Br]PF <sub>6</sub> .	23
1.9	The ‘atypical’ spectrum of SeCNcobalamin with its reduced $\gamma$ band occurring at 371 nm along with other bands, of greater intensity, in the region of 300–350 nm.	24
1.10	Organocobalamins with different R groups most probably occur in solution as a base-on form (A) and two base-off forms (B and C), which are either five- or six-coordinate.	37
2.1	Stopped-flow spectrometer used to collect all the kinetic data in this work.	50
2.2	An example of a trace recorded for the OCN <sup>-</sup> ligand with the fit obtained using a non-linear least-squares program employing a Newton–Raphson procedure. The yellow trace represents a plot of the residuals.	51



**List of Figures** (*continued*)

2.3	HPLC chromatogram of aqua-10-nitrosocobalamin (retention time 7.17 min) using the gradient elution programme reported in Table 2.2.	55
2.4	Crystals of aqua-10-nitrosocobalamin grown from a 50% acetonitrile/water solution and obtained by slow evaporation of the solution.	57
2.5	HPLC chromatogram of aqua-10-aminocobalamin (retention time 6.57 min) using the gradient elution programme reported in Table 2.2.	58
3.1	Spectrophotometric titration of an aquacobalamin solution in a 2.00 cm pathlength cell at the temperatures 5.0 (green), 15.0 (black), 25.0 (blue) and 35.0 °C (red), 0.5 M ionic strength. The solid lines are non-linear least-squares fits of Equation 3.1 to the experimental data.	63
3.2	Temperature dependence of $K_a$ for the self-ionisation of the water ligand in aquacobalamin for the measurements performed at 350nm.	64
3.3	Spectrophotometric titration of an aqua-10-chlorocobalamin solution in a 2.00 cm pathlength cell at 25.0 °C, 0.5 M ionic strength. The solid line is a non-linear least squares fit to the experimental data.	65
3.4	Plot of $\ln K_a$ against $1/T$ for the values obtained at 355 nm in order to obtain the activation parameters.	65
3.5	Spectrophotometric titration of an aqua-10-nitrosocobalamin solution in a 2.00 cm pathlength cell at 25 °C, 0.5 M ionic strength.	66
3.6	Temperature dependence of $K_a$ for the self-ionisation of the water ligand in aqua-10-nitrosocobalamin.	67

**List of Figures** (*continued*)

3.7	Spectrophotometric titration of an aquacobalamin solution in 70% ethanol performed in a 2.00 cm pathlength cell at 25 °C and 522 nm. The solid line is a non-linear least-squares fit to the experimental data.	68
3.8	A plot of the partial charge on oxygen (determined from the ZINDO/1 model) for aquacobalamin, aqua-10-chlorocobalamin and aqua-10-nitrosocobalamin against the $pK_a$ .	70
4.1	Plot of the ligand concentration against the observed rate constant, $k_{obs}$ , from the data in Table 4.1, for the reaction of aquacobalamin with some anionic ligands (25 °C, $I = 2.20 \text{ mol dm}^{-3}$ , KCl). Slight saturation behaviour can be seen.	77
4.2	Dependence of the pseudo-first-order rate constant on ligand concentration for all the ligands at 25 °C.	80
4.3	Dependence of the corrected rate constant on ligand concentration and temperature for the substitution of water by $\text{NO}_2^-$ .	81
4.4	Dependence of the second-order rate constant, $k_{II}$ , on the temperature for the substitution of water by $\text{NO}_2^-$ .	81
4.5	The correlation between $\Delta H_{k_4}^\ddagger$ and $\Delta S_{k_4}^\ddagger$ for the saturating rate constant for the substitution of $\text{H}_2\text{O}$ in aquacobalamin by histamine (1), $\text{N}_3^-$ (2), pyridine (3), $\text{SCN}^-$ (4), 4-methylpyridine (5), imidazole (6), $\text{NO}_2^-$ (7), methyl glycinate (8), $\text{S}_2\text{O}_3^{2-}$ (9), $\text{I}^-$ (10) and $\text{NH}_2\text{OH}$ (11).	82
4.6	The correlation between $\Delta H_{k_{II}}^\ddagger$ and $\Delta S_{k_{II}}^\ddagger$ for the second-order rate constant for the substitution of $\text{H}_2\text{O}$ in aquacobalamin by $\text{S}_2\text{O}_3^{2-}$ (1), $\text{NCO}^-$ (2), $\text{SeCN}^-$ (3), $\text{NO}_2^-$ (4), $\text{SCN}^-$ (5) and $\text{N}_3^-$ (6).	84

**List of Figures** (*continued*)

4.7	Plot of $\ln K_{OS}$ against $T^{-1}$ for the ligand hydroxylamine in order to obtain the activation parameters $\Delta H_K$ and $\Delta S_K$ .	84
4.8	The correlation between $\Delta H_K$ and $\Delta S_K$ , calculated from previously obtained values of $K$ (Figure 4.7) for the substitution of $H_2O$ in aquacobalamin by ligands from Ref. 7 ( $\Delta$ ) and Ref. 8 ( $\circ$ ).	88
4.9	The inverse correlation between the enthalpy change, $\Delta H_K$ , for the formation of the precursor complex and the enthalpy of activation, $\Delta H_{k_4}^\ddagger$ , for the substitution of $H_2O$ in aquacobalamin by the ligands $S_2O_3^{2-}$ , $NO_2^-$ , $SCN^-$ and $N_3^-$ . The same relationship can be seen between $\Delta S_K$ and $\Delta S_{k_4}^\ddagger$ in the inset.	90
4.10	Correlation between $\Delta H_{k_{II}}^\ddagger$ , the Mulliken population of the donor atom as well as the energy of the highest occupied orbital of the ligand with $\sigma$ symmetry.	94
4.11	Correlation between $\Delta S_{k_{II}}^\ddagger$ , the Mulliken population of the donor atom as well as the energy of the highest occupied orbital of the ligand with $\sigma$ symmetry.	94
5.1	Aquacobalamin in 70% ethanol and in water. A slightly more concentrated solution of aquacobalamin in 70% ethanol was used to offset the spectra. There is no evidence for binding of ethanol to aquacobalamin as the band positions in the two solutions are identical.	107
5.2	Plot of the corrected rate constant as a function of ligand concentration for the reaction of aquacobalamin with pyridine in 70% ethanol at 25.0 °C. Equation 5.2 was used to fit the data. The dotted line is a straight line fit of the limiting slope.	109

**List of Figures** (*continued*)

5.3	Plot of the corrected rate constant as a function of ligand concentration for the reaction of aquacobalamin with pyridine in water at 25.0 °C. Equation 5.2 was used to fit the data.	109
5.4	Temperature dependence of $k_4$ for the reactions of aquacobalamin with pyridine in both water and 70% ethanol.	114
6.1	The spectral changes showing aquacobalamin (pH = 8.519; 25 °C); followed by addition of iodide to form iodocobalamin (97% $I^-$ binding); and finally, replacement of the axial iodide ligand with imidazole (0.011 g).	120
6.2	Plot of the corrected rate constant as a function of ligand concentration at 25.0 °C for the replacement of iodide in iodocobalamin by imidazole.	121
6.3	Plot of the corrected rate constant as a function of ligand concentration at 24.9 °C for the replacement of iodide in iodocobalamin by azide.	122
6.4	Plot of $1/k_{\text{cor}}$ against $1/[N_3^-]$ at 24.9 °C extrapolates to $1/k_{\text{sat}}$ , the reciprocal of the saturating rate constant, and the ratio of the intercept and slope gives the equilibrium constant, $K_{\text{OS}}$ .	123
6.5	Plot of the corrected rate constant as a function of ligand concentration at 25.0 °C for the replacement of iodide in iodocobalamin by thiosulfate.	125
6.6	Temperature dependence of $k_{\text{II}}$ for the reaction of iodocobalamin with the thiosulfate ligand.	126
6.7	Plot of $\ln K_{\text{OS}}$ against $1/T$ for the ligand thiosulfate in order to obtain an estimate of the values of $K_{\text{OS}}$ for Equation 6.3.	129
6.8	Plot of $\ln K_{\text{OS}}$ against $1/T$ for the ligand iodide in order to obtain an estimate of the values of $K_{\text{OS}}$ for Equation 6.4.	129
7.1	Chloramine-T.	134

**List of Figures** (*continued*)

7.2	Lactone formation on the addition of one mole of chloramine-T at pH 4.0.	134
7.3	X = Me, CN <sup>-</sup> , and H <sub>2</sub> O, respectively, for methyl-10-chlorocobalamin, cyano-10-chlorocobalamin and aqua-10-chlorocobalamin perchlorate. The Cl atom is shown at the 10-position.	137
8.1	Absorption spectra of aqua-10-chlorocobalamin (a) in an aqueous unbuffered neutral solution at 25 °C, 1 cm pathlength cell, and in the presence of: (b) N <sub>3</sub> <sup>-</sup> (0.040 M), (c) OCN <sup>-</sup> (0.17 M), (d) NO <sub>2</sub> <sup>-</sup> (0.010 M), (e) pyridine (0.41 M), (f) SCN <sup>-</sup> (0.11 M), (g) S <sub>2</sub> O <sub>3</sub> <sup>2-</sup> (0.14 M), (h) SeCN <sup>-</sup> (0.084 M), (i) CH <sub>3</sub> NH <sub>2</sub> (0.12 M) and (j) imidazole (0.054 M).	146
8.2	Relationship between the $\gamma$ bands of both Lcobalamin and L-10-chlorocobalamin ( $R^2 = 0.95$ ).	148
8.3	Plot of the titration data for the binding of the ligand SeCN <sup>-</sup> to aquacobalamin at 350 nm (25 °C, $\mu = 2.2$ M).	151
8.4	Plot of the titration data for the binding of the ligand SeCN <sup>-</sup> to aqua-10-chlorocobalamin at 350 nm (25 °C, $\mu = 2.2$ M).	151
8.5	Plot of the residuals for the data of the binding of aqua-10-chlorocobalamin with the ligand SeCN <sup>-</sup> .	152
9.1	The spectra of aqua-10-chlorocobalamin ( $\approx 20$ $\mu$ M) (where water is the axial ligand) and the substitution of the water ligand for azide (0.17 M) and pyridine (0.41 mM).	164
9.2	Plot of the corrected rate constant as a function of ligand concentration, at 25.0 °C, for the reactions of both aquacobalamin and aqua-10-chlorocobalamin with pyridine.	166
9.3	The temperature dependence of $k_4$ for the reaction of pyridine with both aquacobalamin and aqua-10-chlorocobalamin.	168

**List of Figures** (*continued*)

9.4	Plot of the corrected rate constant as a function of ligand concentration, at 25.0 °C, for the reactions of both aquacobalamin and aqua-10-chlorocobalamin with azide.	170
9.5	The temperature dependence of $k_{II}$ for the reaction of azide with both aquacobalamin and aqua-10-chlorocobalamin.	172
9.6	Plot of $\ln K_{OS}$ against $1/T$ for the ligand azide in order to obtain the activation parameters $\Delta H_{K_{OS}}$ and $\Delta S_{K_{OS}}$ .	174
9.7	A plot of the partial charge on cobalt (determined from an MPA using semi-empirical MO calculations and the ZINDO/1 model) for both (●) aquacobalamin and (▲) aqua-10-chlorocobalamin against the Co–O bond length.	179
9.8	A plot of the partial charge on oxygen (determined from an MPA using semi-empirical MO calculations and the ZINDO/1 model) for both (+) aquacobalamin and (▲) aqua-10-chlorocobalamin against the Co–O bond length.	179
9.9	The compensation effect for the reaction between $\Delta H^\ddagger$ and $\Delta S^\ddagger$ for the reactions between pyridine ( $k_4$ and $k_{-4}$ ) and azide ( $k_4$ ) for both aqua-10-chlorocobalamin and aquacobalamin.	181
9.10	A plot of the corrected rate constants against ligand concentration for the reactions of pyridine and azide with aqua-10-chlorocobalamin, at 25 °C, in order to show the difference in their saturating rates.	182
10.1	Aqua-10-nitrosocobalamin, where the H atom in the 10-position has been replaced with a NO group.	186

**List of Figures** (*continued*)

10.2	The UV-vis spectra of: (a) 119 $\mu\text{M}$ solution of aqua-10-nitrosocobalamin, pH 7, in aqueous solution; (b) Addition of one equivalent of $\text{CN}^-$ leads to the slow conversion to cyano-10-nitrosocobalamin; (c) Excess $\text{CN}^-$ results in the formation of dicyano-10-nitrosocobalamin; and (d) At pH > 13 this compound is converted into the isonitroso derivative.	188
10.3	The significant differences between $^{13}\text{C}$ NMR resonances of aquacobalamin and aqua-10-chlorocobalamin where the values are from $\delta_{10\text{-NO-H}_2\text{OCbl}^+} - \delta_{\text{H}_2\text{OCbl}^+}$ .	191
10.4	A comparison of the spectrum of aquacobalamin with those of aqua-10-chlorocobalamin and aqua-10-nitrosocobalamin, where Cl and NO have substituted the H at the 10-position, respectively.	192
10.5	The addition of 1.2 M pyridine to aqua-10-nitrosocobalamin after 72 hours, indicating that no ligand substitution of $\text{H}_2\text{O}$ has occurred.	193
10.6	Photographs of aqua-10-nitrosocobalamin (a) and (b) and cobalamin <sup>4</sup> (c) and (d) crystals. These show that aqua-10-nitrosocobalamin crystals are much thinner and plate-like when compared with the prismic shaped cobalamin crystals. This has serious consequences on the diffracting ability of the aqua-10-nitrosocobalamin crystals, which makes X-ray diffraction studies by conventional methods extremely difficult.	195

**List of Figures** (*continued*)

10.7	Photographs of the crystals obtained from the mixture of aqua-10-nitrosocobalamin and 5–10% aquacobalamin using the hanging drop method.	196
11.1	The spectra of aqua-10-aminocobalamin and dicyano-10-aminocobalamin determined in aqueous solution.	200
11.2	Decomposition products of aqua-10-aminocobalamin. The blue line shows the compound after exposure to light with the initial disappearance of the $\alpha\beta$ bands and the red line is the formation of a stable yellow corrinoid after heating the compound.	200
11.3	Absorption spectra of aqua-10-aminocobalamin (a) in an aqueous unbuffered neutral solution at 25 °C, 1 cm pathlength cell, and in the presence of (b) pyridine (0.41 M), (c) $\text{SCN}^-$ (0.060 M), (d) $\text{S}_2\text{O}_3^{2-}$ (0.24 M), (e) $\text{NO}_2^-$ (0.0072 M), (f) $\text{CH}_3\text{NH}_2$ (1.25 M), (g) $\text{N}_3^-$ (0.072 M), (h) $\text{OCN}^-$ (0.030 M), (i) $\text{SeCN}^-$ (0.50 M).	201



**LIST OF TABLES**

<b>Table</b>		<b>Page</b>
1.1	Some naturally occurring nucleotide analogues of vitamin B <sub>12</sub> .	5
1.2	Rate constants for the ligand substitution of water in various Co <sup>III</sup> (N <sub>4</sub> ) complexes.	14
1.3	Correlation between absorption spectra and NMR spectra of cobalamins and cobinamides.	15
1.4	Ground-state <i>cis</i> - and <i>trans</i> -effects on the bond lengths of some Co(III) complexes.	17
1.5	Ground-state <i>cis</i> - and <i>trans</i> -effects on CN stretching frequencies in Co(III) complexes.	18
1.6	The position of the $\gamma$ band with the replacement of the water ligand by various ligands indicating the steric affect that the ligands have on the cobalt atom.	26
1.7	Ligands that have been used for ligand substitution reaction studies with aquacobalamin.	34
2.1	Buffers used.	47
2.2	HPLC elution programme used throughout this work.	55
3.1	Thermodynamic parameters from the ionisation of the water ligand in aquacobalamin, aqua-10-chlorocobalamin and aqua-10-nitrosocobalamin.	69
4.1	Rate constants and activation parameters for the reaction of aquacobalamin with various ligands.	75
4.2	Rate constants and activation parameters for the reaction of aquacobalamin with various ligands ( $I = 2.2 \text{ mol dm}^{-3}$ , NaNO <sub>3</sub> ).	79
4.3	Thermodynamic parameters for the formation of the outer-sphere complex between aquacobalamin and an incoming ligand.	85

**List of Tables** (*continued*)

4.4	Mulliken populations of the donor atoms as well as the symmetry and energy of the highest occupied MO with $\sigma$ symmetry of the ligands. Calculations were performed by an <i>ab initio</i> molecular orbital method using a minimal STO-6G* basis set.	92
4.5	Correlation statistics for the Mulliken population of the donor atom and the energy of the highest occupied molecular orbital with $\sigma$ symmetry with $\Delta H_{k_{II}}^\ddagger$ and $\Delta S_{k_{II}}^\ddagger$ , respectively.	95
5.1	Saturating rate constants and pre-equilibrium values obtained for the reactions of aquacobalamin with pyridine in both water (using Equation 5.2) and 70% ethanol (using Equations 5.2, 5.3 and 5.4).	111
5.2	Rate constants for the comparison of the reaction of aquacobalamin with thiourea in both 70% acetonitrile and 70% dioxane; aquacobalamin with thiocyanate in acetonitrile; and aquacobalamin with pyridine in 70% ethanol.	112
5.3	Activation parameters for $k_4$ determined from a plot of $\ln(k_4h/k_B T)$ against $1/T$ for the reactions of aqua-cobalamin with pyridine in both water and 70% ethanol.	114
6.1	Rate constants, $k_4$ , and outer-sphere formation constants, $K_{OS}$ , for the replacement of iodide in iodocobalamin with imidazole and azide, and the second order rate constant for the reaction with thiosulfate.	124
6.2	Equilibrium constants for the coordination of the ligands thiosulfate, iodide and azide by aquacobalamin, along with the thermodynamic parameters for the formation of an outer-sphere complex between the incoming ligand and aquacobalamin calculated from these equilibrium constants.	128

**List of Tables** (*continued*)

6.3	Calculated values of $\Delta H_{k_4}^\ddagger$ and $\Delta S_{k_4}^\ddagger$ for the reactions of both aquacobalamin and iodocobalamin with the ligands $S_2O_3^{2-}$ and $N_3^-$ .	131
6.4	A comparison between the $k_{II}$ values and resulting activation parameters of the reaction of both aquacobalamin and iodocobalamin with thiosulfate, respectively.	131
7.1	The effect on the electronic spectrum of cyanocobalamin on changing the atom in the C10 position.	136
7.2	Bond lengths and bond angles for the crystal structure of XCbl and X-10-ClCbl where X = Me, $CN^-$ and $H_2O$ .	139
8.1	Some formation constants, reproduced from Pratt, <sup>15</sup> for the substitution of water by cyanide, methylisocyanide, azide, hydroxide and imidazole, with a variety of ligands in the <i>trans</i> -position.	143
8.2	A comparison of the effect of the axial ligand on the wavelength of the $\gamma$ , $\beta$ and $\alpha$ bands for aquacobalamin and aqua-10-chlorocobalamin.	147
8.3	A comparison of the $\log K$ values, determined in this experiment and those reported elsewhere, obtained from the binding of various ligands to aquacobalamin in the axial position.	153
8.4	A comparison of the $\log K$ values obtained from the binding of various ligands in the axial position to both aquacobalamin and aqua-10-chlorocobalamin.	154
8.5	Partial charges on the chlorine and cobalt atoms determined by Brown <i>et al.</i> <sup>24</sup> for MeCbl, CNCbl and $H_2OCbl^+$ as well as their 10-Cl analogues.	155
9.1	Ground state <i>cis</i> -effects in Co(III) systems.	162

**List of Tables** (*continued*)

9.2	Rate and equilibrium constants for the coordination of pyridine by aquacobalamin and aqua-10-chlorocobalamin.	165
9.3	A comparison of the corrected rate constants obtained from the reaction of pyridine with aquacobalamin and aqua-10-chlorocobalamin at 25 °C.	167
9.4	Activation parameters for the reaction of pyridine with aquacobalamin and aqua-10-chlorocobalamin.	168
9.5	A comparison of the rate constants obtained from the reaction of azide with aquacobalamin and aqua-10-chlorocobalamin at 15.0 °C.	171
9.6	Second-order rate constants, $k_{II}$ , and corresponding activation parameters for the reaction of aquacobalamin and aqua-10-chlorocobalamin with azide.	172
9.7	Equilibrium constants for the coordination of azide by aquacobalamin.	173
9.8	Calculated parameters of $\Delta H_{k_4}^\ddagger$ and $\Delta S_{k_4}^\ddagger$ using previously determined $K_{OS}$ values (Table 9.7) and the experimentally determined activation parameters $\Delta H_{k_{II}}^\ddagger$ and $\Delta S_{k_{II}}^\ddagger$ .	175
9.9	Calculated $k_4$ values using the Fuoss-Eigen equation, assuming a spherical aquacobalamin with a net charge of +2 and +1, respectively.	176
9.10	Partial charges on cobalt and oxygen (determined from an MPA using semi-empirical MO calculations and the ZINDO/1 model) for both aquacobalamin and aqua-10-chlorocobalamin.	178
9.11	Summary of the activation parameters for the reaction of aquacobalamin and aqua-10-chlorocobalamin with both azide and pyridine.	180

**List of Tables** (*continued*)

10.1	A comparison of the band peaks obtained for some of the nitroso derivatives.	189
10.2	The significant differences in the $^{13}\text{C}$ NMR resonances, in ppm, between aqua-10-nitrosocobalamin and aquacobalamin, and between aqua-10-chlorocobalamin and aquacobalamin.	190
11.1	A comparison of the effect of the axial ligand on the wavelength of the $\gamma$ , $\beta$ and $\alpha$ bands for aquacobalamin and aqua-10-chlorocobalamin.	202

**LIST OF ABBREVIATIONS**

CSD	Cambridge structural database
DH	monoanion of dimethylglyoxime
dmbzim	5,6-dimethylbenzimidazole
ESR	electron spin resonance
En	ethylenediamine
FAB(MS)	fast atom bombardment mass spectrometry
HMBC	heteronuclear multiple bond correlation
HMQC	<sup>1</sup> H-detected heteronuclear multiquantum correlation
HPLC	high pressure liquid chromatography
MPA	Mulliken population analysis
NMR	nuclear magnetic resonance
PM3	a reparameterisation of the Austin model 1 force field
ROESY	rotating frame Overhauser effect spectroscopy
Saloph	salicylaldehydophenylenediimine
TOCSY	total correlation spectroscopy
UV-vis	ultraviolet-visible
ZINDO/1	Zerner's non-spectroscopic version of the intermediate neglect of differential overlap

**Buffers**

CAPS	3-[cyclohexylamino]-1-propanesulfonic acid <sup>†</sup>
CHES	2-[N-cyclohexylamino]-ethanesulfonic acid
HEPES	N-[2-Hydroxyethyl]piperazine-N'[2ethanesulfonic acid]
MES	2-[N-Morpholino]ethanesulfonic acid
MOPS	3-[N-Morpholino]propanesulfonic acid
Tris	Tris(hydroxymethyl)aminomethane

<sup>†</sup> Sulfur rather than sulphur is the IUPAC-recommended spelling and will be used throughout this thesis for related words.

**Symbols**

$A$	Absorbance
$j$	coupling constant
$K_a$	ionisation constant
$K_L$	acid dissociation constant for L
$K_{OS}$	Equilibrium constant for formation of the outer-sphere complex between L and cobalamin under investigation
$K_{obs}$	experimentally determined equilibrium constant for coordination of L by the cobalamin under investigation
$k_{cor}$	observed rate constant corrected for pH effects
$k_{obs}$	experimentally determined rate constant for the substitution of H <sub>2</sub> O by L in the cobalamins
$k_{sat}$	saturation rate constant